

ORIGINAL RESEARCH ORJİNAL ARAŞTIRMA

DOI: 10.5336/healthsci.2025-108556

Relationship Between Adherence to Cardioprotective Diet Recommendations and Disease Risk Awareness: A Cross-Sectional Study

Kardiyoprotektif Diyet Önerilerine Uyum ile Hastalık Riski Farkındalığı Arasındaki İlişki: Kesitsel Araştırma

¹Büşra BAŞAR GÖKCEN^a, ²Dila ÖZDEMİR^a, ³Sena AKSU^b

^aMuğla Sıtkı Koçman University Fethiye Faculty of Health Sciences, Department of Nutrition and Dietetics, Muğla, Türkiye

^bGazi University Faculty of Health Sciences, Department of Nutrition and Dietetics, Ankara, Türkiye

ABSTRACT Objective: This study aims to assess the alignment of the Mediterranean diet with the 2021 American Heart Association (AHA) Dietary Guidelines and examine the relationship between adherence to cardioprotective dietary recommendations and awareness of cardiovascular disease (CVD) risk factors. **Material and Methods:** This study included 200 individuals diagnosed with CVD who applied to the Cardiology Clinic of Fethiye State Hospital. Data were collected through face-to-face surveys, assessing participants' sociodemographic characteristics, body mass index (BMI) and blood pressure values, and dietary habits. Adherence to cardioprotective dietary recommendations was evaluated using the Mediterranean Diet Adherence Scale (MEDAS) and a 10-item questionnaire assessing adherence with the 2021 AHA Dietary Guidelines. Additionally, CVD risk awareness was assessed using the Cardiovascular Disease Risk Awareness Assessment Questionnaire (CDRAAQ). **Results:** Participants with higher adherence to the Mediterranean diet had significantly greater awareness of CVD risk factors ($p<0.05$) compared to those with low or moderate adherence. They were also older ($p=0.010$) and had lower diastolic blood pressure ($p=0.033$), while no significant differences were observed in BMI or systolic blood pressure ($p>0.05$). Individuals who followed the AHA dietary guidelines demonstrated higher Mediterranean diet adherence and greater awareness of CVD risk factors ($p<0.05$). Additionally, CDRAAQ scores were positively correlated with MEDAS scores and negatively correlated with BMI and blood pressure ($p<0.05$). **Conclusion:** These findings emphasize the role of disease-specific awareness and educational interventions in promoting adherence to cardioprotective dietary patterns among individuals with CVD. Enhancing awareness of CVD risk factors may contribute to improved dietary adherence and cardiovascular health outcomes.

ÖZET Amaç: Bu çalışma, Akdeniz diyetinin 2021 Amerikan Kalp Derneği'nin [American Heart Association (AHA)] Beslenme Kılavuzu ile uyumunu değerlendirmeyi ve kardiyoprotektif beslenme önerilerine uyum ile kardiyovasküler hastalık (KVH) risk faktörleri farkındalığı arasındaki ilişkiyi incelemeyi amaçlamaktadır. **Gereç ve Yöntemler:** Çalışmaya, Fethiye Devlet Hastanesi Kardiyoloji Kliniği'ne başvuran ve KVH tanısı almış 200 birey dâhil edilmiştir. Veriler, yüz yüze uygulanan anketler yoluyla toplanmış; katılımcıların sosyodemografik özellikleri, beden kitle indeksi (BKİ), kan basıncı değerleri ve beslenme alışkanlıkları değerlendirilmiştir. Kardiyoprotektif beslenme önerilerine uyum, Akdeniz Diyeti Uyum Ölçeği [Mediterranean Diet Adherence Scale (MEDAS)] ve 2021 AHA Beslenme Kılavuzu'na uyumu değerlendiren 10 maddelik bir anket ile ölçülmüştür. Ayrıca KVH risk farkındalığı, Kardiyovasküler Hastalık Risk Farkındalığı Değerlendirme Anketi [Cardiovascular Disease Risk Awareness Assessment Questionnaire (CDRAAQ)] kullanılarak değerlendirilmiştir. **Bulgular:** Akdeniz diyetine yüksek uyum gösteren katılımcılar, düşük veya orta uyum gösterenlere kıyasla KVH risk faktörleri farkındalığında anlamlı derecede daha yüksek puan almıştır ($p<0.05$). Ayrıca, bu katılımcılar daha yaşlı ($p=0.010$) olup, daha düşük diyastolik kan basıncına sahipti ($p=0.033$). Ancak BKİ ve sistolik kan basıncı açısından anlamlı bir fark bulunmamıştır ($p>0.05$). AHA beslenme kılavuzlarına uyan bireylerin Akdeniz diyetine daha yüksek uyum gösterdiği ve KVH risk faktörleri farkındalıklarının daha yüksek olduğu belirlenmiştir ($p<0.05$). Ek olarak, CDRAAQ puanları MEDAS puanları ile pozitif, BKİ ve kan basıncı ile negatif korelasyon göstermiştir ($p<0.05$). **Sonuç:** Bu bulgular, KVH'ye özgü farkındalık ve eğitim müdahalelerinin, kardiyoprotektif beslenme kalıplarına uyumu teşvik etmede önemli bir role sahip olduğunu vurgulamaktadır. KVH risk faktörleri farkındalığının artırılması, beslenme uyumunun ve kardiyovasküler sağlık sonuçlarının iyileştirilmesine katkıda bulunabilir.

Keywords: Mediterranean diet; cardiovascular disease; risk factors

Anahtar Kelimeler: Akdeniz diyeti; kardiyovasküler hastalık; risk faktörleri

TO CITE THIS ARTICLE:

Başar Gökcen B, Özdemir D, Aksu S. Relationship between adherence to cardioprotective diet recommendations and disease risk awareness: A cross-sectional study. Türkiye Klinikleri J Health Sci. 2025;10(3):550-60.

Correspondence: Büşra BAŞAR GÖKCEN

Muğla Sıtkı Koçman University Fethiye Faculty of Health Sciences, Department of Nutrition and Dietetics, Muğla, Türkiye

E-mail: busrabasar@mu.edu.tr



Peer review under responsibility of Türkiye Klinikleri Journal of Health Sciences.

Received: 15 Jan 2025

Received in revised form: 10 Mar 2025

Accepted: 24 Mar 2025

Available online: 30 Jun 2025

2536-4391 / Copyright © 2025 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Cardiovascular diseases (CVDs) remain a leading cause of premature death and disability globally, driven by environmental, lifestyle, behavioral, and genetic factors.^{1,2} A balanced diet, physical activity, smoking cessation, and other modifiable factors such as moderate alcohol intake and healthy weight contribute to CVD prevention.³

According to American and European guidelines, diet plays a crucial role in influencing cardiovascular risk factors and is central to disease prevention. Both guidelines promote the intake of whole foods, replacing saturated with unsaturated fats, limiting trans fats and added sugars, and reducing sodium to support cardiovascular health. Additionally, they recommend the Dietary Approaches to Stop Hypertension (DASH) and Mediterranean diets, both of which encourage the consumption of fruits, vegetables, whole grains, legumes, and nuts to support healthy eating habits.⁴⁻⁶ Although there is a general consensus on dietary patterns proven to be effective in reducing cardiovascular risk, knowledge gaps still remain.³

A recent study evaluated the alignment of popular diets with the 2021 American Heart Association (AHA) Dietary Guidelines, categorizing dietary patterns into four groups based on their adherence levels. The results indicated that DASH, Mediterranean, Pescetarian, and Vegetarian diets demonstrated the strongest alignment, whereas Vegan, low-fat, and low-carbohydrate diets ranked at lower adherence levels.⁷

As the prevalence of CVD risk factors continues to rise worldwide, it is crucial to focus on understanding the causes of this disease and increasing awareness. Although symptoms are one of the primary reasons patients seek medical care for chronic diseases, they may sometimes go unnoticed or be disregarded until the disease reaches an advanced stage. At this point, disease awareness plays a critical role in effective management. Awareness of CVD and its associated risk factors is anticipated to be highly significant in encouraging individuals to adopt healthy lifestyle changes and adhere to disease-specific dietary recommendations.^{8,9}

While existing guidelines provide dietary recommendations to protect cardiovascular health, the extent to which individuals adhere to these recommendations and the role of awareness in this process remain unclear. This study aims to evaluate the alignment of the Mediterranean diet with the 2021 AHA Dietary Guidelines and analyze the relationship between adherence to cardioprotective dietary recommendations and awareness of CVD risk factors, thereby addressing the knowledge gap in this field.

MATERIAL AND METHODS

SAMPLE OF THE STUDY

The population of this study consists of individuals diagnosed with CVD who applied to the Cardiology Clinic of Fethiye State Hospital under the Muğla Provincial Health Directorate. Individuals diagnosed with CVD who met the inclusion criteria, participated voluntarily between October–December 2022, and completed the survey were included (n=200). The inclusion criteria for the study were being over 18 years old, voluntarily participating in the study, and having no verbal communication barriers. The exclusion criteria included being pregnant or in the breastfeeding period, as well as being physically, psychologically, or socially unable to complete the survey. The sample size was calculated using the Power Analysis Program (G-Power 3.1.9.7; Heinrich Heine University, Düsseldorf, Germany), and a one-way ANOVA variance analysis was performed. In the calculation, a medium effect size ($f=0.25$), a 5% margin of error ($\alpha=0.05$), and 80% power ($1-\beta=0.80$) were considered, resulting in a required sample size of 195.

This study was approved by the Muğla Sıtkı Koçman University Ethics Committee (June 29, 2022; No. 220090-89) and the Muğla Provincial Health Directorate (October 14, 2022).

Written informed consent was obtained from all participants. Participants were provided with preliminary information regarding the procedures to be carried out within the scope of the study and their expected involvement. If they voluntarily agreed to participate, their consent was obtained through the Informed Voluntary Consent Form.

DATA COLLECTION TOOLS

The questionnaire was administered by the researcher using the face-to-face interview method to collect data. The questionnaire assessed the sociodemographic characteristics and lifestyle choices of the individuals. Additionally, individuals' body weight and height were recorded based on their self-reported measurements. Individuals' usual systolic blood pressure (SBP) and diastolic blood pressure (DBP) levels were also recorded based on their self-reported measurements.

The Cardiovascular Disease Risk Awareness Assessment Questionnaire (CDRAAQ) was used to assess awareness of CVD risk, while the Mediterranean Diet Adherence Scale (MEDAS) measured adherence to the Mediterranean diet. Additionally, adherence to the recommendations outlined in the AHA 2021 Dietary Guidelines for Improving Cardiovascular Health was evaluated.

Cardiovascular Disease Risk Awareness Assessment Questionnaire

The first version of the CDRAAQ developed by the European Society of Cardiology consisted of 65 items. Later, Woringer et al. adapted this form to the British society in 2017 and obtained a 26-item scale.¹⁰ Vural Doğru et al. adapted this version to the Turkish society in 2021 and obtained its validity and reliability.¹¹

The first 8 items in the scale are knowledge questions, with responses given as “true”, “false”, or “I don't know”. Each correct answer is scored as 1 point, while incorrect or “I don't know” responses are scored as 0 points. The correct answer for the first 7 questions is “true”, while the correct answer for question 8 is “false”. Apart from this, the scale consists of three sub-dimensions: perceived heart attack/stroke risk (7 items), healthy eating intentions (5 items), and perceived benefits and intentions for change (2 items). The scale is structured as a 4-point Likert scale, where “1=Strongly Disagree” and “4=Strongly Agree”. Additionally, items 15-16 are reverse-scored.¹¹

Higher scores in the sub-dimensions indicate greater perceived CVD risk, stronger intentions for

healthy eating, and readiness for behavioral change. Increased total and subscale scores reflect greater cardiovascular disease awareness.

In the original version of the scale, the internal consistency reliability of the factor structure was evaluated using Cronbach's α coefficient. The Perceived Risk of Heart Attack/Stroke factor had $\alpha=0.85$, the Perceived Benefits and Intentions to Change factor had $\alpha=0.82$, and the Healthy Eating Intentions factor had $\alpha=0.56$. In the Turkish version, these values were found to be $\alpha=0.915$, $\alpha=0.792$, and $\alpha=0.682$, respectively.^{10,11}

Mediterranean Diet Adherence Scale

MEDAS was used to assess participants' adherence with the Mediterranean diet. This scale was developed by Martínez-González et al. in 2012, and its Turkish validity and reliability study was conducted by Özkan Pehlivanoglu et al. in 2020. The scale consists of 14 questions, with 1 or 0 points assigned to each question based on consumption amounts, and the total score range for MEDAS has been specified as 0-14. These scores are summed to obtain a total score, with adherence categorized as low (≤ 5 points), medium (6-9 points), or high (≥ 10 points).^{12,13}

The Turkish version of MEDAS demonstrated good internal consistency (Cronbach's $\alpha = 0.829$).^{12,13}

Assessment of adherence to AHA Dietary Guidelines

In the AHA 2021 Guidelines for Improving Cardiovascular Health, 10 key recommendations were outlined, 5 of which align with the principles of the Mediterranean diet. AHA recommendations were evaluated using yes/no items. Participants were encouraged to consume diverse fruits and vegetables, choose whole grains, prefer healthy protein sources, use plant-based oils, and limit ultra-processed foods.⁶

STATISTICAL ANALYSIS

All statistical analyses were performed using the SPSS Statistics Version 24.0 (SPSS, Inc.; Chicago, USA). Prior to conducting the analyses, the normality of continuous variables was assessed using the Kolmogorov-Smirnov test, as it is more appropriate for larger sample sizes ($n > 50$ per group). Addition-

ally, visual methods, including histograms, Q-Q plots, and box plots, were examined to further support the normality assessment. Skewness and Kurtosis values were also considered, with values within the range of ± 1 indicating an approximately normal distribution. For descriptive statistics, frequency and percentage were calculated for qualitative variables, while mean and standard deviation were reported for quantitative variables.

For group comparisons, parametric tests were used for normally distributed variables, specifically the independent samples t-test for 2 independent groups. When the normality assumption was not met, non-parametric tests were applied, namely the Mann-Whitney U test for comparisons between 2 independent groups. For categorical variables, group comparisons were performed using the chi-square test.

Correlation analyses were conducted using Pearson correlation for normally distributed variables and Spearman correlation for non-normally distributed variables.

A $p < 0.05$ was considered statistically significant.

RESULTS

A total of 200 patients diagnosed with CVD participated in this study. Participants had a mean age of 60.0 ± 13.4 years and BMI of 29.6 ± 5.4 kg/m²; 48.5% were female, 75% married, 83% unemployed, and 48.5% had primary education. Additionally, 25% of the participants reported smoking, while 17.5% reported alcohol consumption. Just over half of the participants (50.5%) had at least one comorbid condition, and 20.5% reported regular use of nutritional supplements.

Table 1 presents CDRAAQ scores and health indicators by Mediterranean diet adherence level. Among the 200 participants, those with high Mediterranean diet adherence had a significantly higher average CDRAAQ total score (45.2 ± 10.8) than the low or medium adherence group (40.8 ± 12.6 , $p < 0.05$). Similar significant results were shown in other sub-factors except for the knowledge sub-factor

($p = 0.456$). High adherence was associated with significantly higher CDRAAQ SF2 ($p = 0.036$), SF3 ($p = 0.010$), and SF4 ($p = 0.049$) scores.

Participants in the high adherence group were significantly older (62.5 ± 11.2 years vs. 56.3 ± 15.6 years, $p = 0.010$) and had significantly lower DBP (76.7 ± 12.3 mmHg vs. 80.5 ± 13.8 mmHg, $p = 0.033$). However, there were no statistically significant differences between the groups in terms of BMI (29.3 ± 5.3 kg/m² vs. 30.1 ± 5.5 kg/m², $p = 0.313$) and SBP (133.3 ± 18.5 mmHg vs. 135.2 ± 22.6 mmHg, $p = 0.570$).

Table 2 shows correlations between Mediterranean diet adherence and CVD risk awareness. MEDAS total score correlated positively with CDRAAQ SF3 ($r = 0.184$, $p < 0.01$) and total awareness ($r = 0.155$, $p < 0.05$). BMI and DBP showed negative correlations with SF4, while SBP was inversely associated with SF3, SF4, and total score ($p < 0.05$).

The alignment of the Mediterranean diet with the 2021 AHA Dietary Guidance is presented in Table 3. Overall, participants who reported maintaining a healthy body weight were more likely to be in the high Mediterranean diet adherence group (68.1%) compared to those in the low or medium adherence group (53.1%) ($p = 0.032$). Dietary behaviors that aligned with AHA recommendations were significantly associated with higher adherence to the Mediterranean diet. Specifically, participants who consumed a wide variety of fruits and vegetables demonstrated significantly higher MEDAS scores than non-adherents ($p = 0.027$). Similarly, adherence to the consumption of fish and seafood was notably stronger among those with higher Mediterranean diet adherence, as reflected by their significantly higher MEDAS scores ($p = 0.002$). In terms of proportional adherence, 80.7% of individuals in the high adherence group complied with the fish and seafood recommendation, compared to 61.7% in the low or medium adherence group ($p = 0.003$).

Preferences for lean protein sources also played a significant role. Participants who selected lean cuts of poultry had higher MEDAS scores than those who did not ($p = 0.017$). Moreover, the use of liquid plant

TABLE 1: CVD risk awareness and selected health indicators in relation to participants' Mediterranean diet adherence levels

Variables	Total (n=200)	Low or medium adherence (n=81)	High adherence (n=119)	p value
Age (years)	60.0±13.4 (61)	56.3±15.6 (57)	62.5±11.2 (62)	0.010
BMI (kg/m ²)	29.6±5.4 (29)	30.1±5.5 (29)	29.3±5.3 (29)	0.313
SBP (mmHg)	134.1±20.2 (130)	135.2±22.6 (130)	133.3±18.5 (130)	0.570
DBP (mmHg)	78.2±13.0 (80)	80.5±13.8 (80)	76.7±12.3 (80)	0.033
CDRAAQ TS	43.4±11.7 (44)	40.8±12.6 (42)	45.2±10.8 (48)	0.011
CDRAAQ SF1	6.3±1.7 (7)	6.1±1.8 (7)	6.4±1.6 (7)	0.456
CDRAAQ SF2	19.6±6.6 (22)	18.5±7.0 (21)	20.3±6.3 (23)	0.036
CDRAAQ SF3	12.6±4.0 (13)	11.6±4.3 (11)	13.2±3.6 (13)	0.010
CDRAAQ SF4	5.0±3.0 (7)	4.6±2.9 (6)	5.3±3.0 (7)	0.049
MEDAS TS	9.9±2.3 (10)	7.6±1.5 (8)	11.4±1.3 (11)	0.000

BMI: Body mass index; SBP: Systolic blood pressure; DBP: Diastolic blood pressure; CDRAAQ: Cardiovascular Disease Risk Awareness Assessment Questionnaire; TS: Total score; SF: Sub-factor; MEDAS: Mediterranean Diet Adherence Scale. CDRAAQ SF1: Knowledge; CDRAAQ SF2: Heart attack/stroke risk; CDRAAQ SF3: Healthy eating intentions; CDRAAQ SF4: Perceived benefits and intentions for change.

This table presents the comparison of Cardiovascular Disease Risk Awareness (CDRAAQ TS and sub-factors) and Selected Health Indicators (BMI, SBP, DBP) based on participants' adherence levels to the Mediterranean diet (MEDAS TS). Values are expressed as mean±standard deviation (median). For normally distributed variables, the independent samples t-test was used, whereas the Mann-Whitney U Test was applied for non-normally distributed variables.

TABLE 2: Correlation between Mediterranean diet adherence, CVD risk awareness, and selected health indicators

Variables	MEDAS TS	BMI (kg/m ²)	DBP (mmHg)	SBP (mmHg)
MEDAS TS	1.000			
BMI (kg/m ²)	-0.051	-		
DBP (mmHg)	-0.136	0.153*	-	
SBP (mmHg)	-0.055	0.150*	0.607**	-
CDRAAQ SF1	0.041	-0.138	-0.014	-0.077
CDRAAQ SF2	0.108	-0.036	-0.039	-0.107
CDRAAQ SF3	0.184**	-0.111	-0.063	-0.147*
CDRAAQ SF4	0.120	-0.220**	-0.159*	-0.295**
CDRAAQ TS	0.155*	-0.130	-0.084	-0.183**

*Indicates significance at the 0.05 level; **indicates significance at the 0.01 level. MEDAS: Mediterranean Diet Adherence Scale; TS: Total score; BMI: Body mass index; DBP: Diastolic blood pressure; SBP: Systolic blood pressure; CDRAAQ: Cardiovascular Disease Risk Awareness Assessment Questionnaire; SF: Sub-factor. CDRAAQ SF1: Knowledge; CDRAAQ SF2: Heart attack/stroke risk; CDRAAQ SF3: Healthy eating intentions; CDRAAQ SF4: Perceived benefits and intentions for change.

The correlations were assessed using the Pearson correlation analysis for normally distributed variables and the Spearman rank correlation analysis for non-normally distributed variables. The table presents correlation coefficients (r) to evaluate the strength and direction of relationships between variables. Statistically significant correlations are indicated by p values, with p<0.05 considered significant and p<0.01 considered highly significant.

oils instead of tropical oils was associated with significantly higher MEDAS scores (p=0.008). This preference was more prominent in the high adherence group, where 84.0% adhered to this recommendation, compared to 71.6% in the low or medium adherence group (p=0.034). Limiting added sugar intake also emerged as a significant factor. Participants who restricted their intake of added sugars had higher

MEDAS scores (p=0.001), with 76.5% of the high adherence group complying with this recommendation, compared to 53.1% in the low or medium adherence group (p=0.001). Lastly, participants who adhered to dietary recommendations regardless of where the food was prepared or consumed also demonstrated significantly higher MEDAS scores (p=0.047).

TABLE 3: Alignment of the Mediterranean diet with the 2021 AHA Dietary Guidance

AHA features	MEDAS TS	Low or medium adherence (n=81)	High adherence (n=119)
(1) Energy balance needed to maintain a healthy weight Adherents (n=124, 62.0%) Non-adherents (n=76, 38.0%) p value	10.1±2.2 (10.0) 9.5±2.4 (9.5) 0.071	43 (53.1%) 38 (46.9%) 0.032	81 (68.1%) 38 (31.9%)
(2) Eat plenty of vegetables and fruits, a wide variety Adherents (n=120, 60.0%) Non-adherents (n=80, 40.0%) p value	10.2±2.2 (10.0) 9.4±2.4 (10.0) 0.027	43 (53.1%) 38 (46.9%) 0.100	77 (64.7%) 42 (35.3%)
(3) Choose mostly whole grains rather than refined grains Adherents (n=128, 64.0%) Non-adherents (n=72, 36.0%) p value	10.0±2.3 (10.0) 9.7±2.3 (10.0) 0.734	54 (66.7%) 27 (33.3%) 0.517	74 (62.1%) 45 (37.8%)
(4) ^a Mostly protein from plants (legumes and nuts) Adherents (n=140, 70.0%) Non-adherents (n=60, 30.0%) p value ^b Fish and seafood Adherents (n=146, 73.0%) Non-adherents (n=54, 27.0%) p value ^c Low-Fat or fat-free dairy products instead of full-fat dairy Adherents (n=40, 20.0%) Non-adherents (n=160, 80.0%) p value ^{d1} If consuming poultry, choose lean cuts Adherents (n=132, 66.0%) Non-adherents (n=68, 34.0%) p value ^{d2} If consuming meat, choose lean cuts Adherents (n=112, 56.0%) Non-adherents (n=88, 44.0%) p value	10.0±2.2 (10.0) 9.5±2.5 (9.5) 0.129 10.2±2.3 (10.0) 9.0±2.3 (9.0) 0.002 9.9±2.8 (10.0) 9.9±2.2 (10.0) 0.901 10.2±2.1 (10.0) 9.3±2.6 (10.0) 0.017 10.1±2.1 (10.0) 9.6±2.5 (10.0) 0.090	51 (63.0%) 30 (37.0%) 0.073 50 (61.7%) 31 (38.3%) 0.003 18 (22.2%) 63 (77.8%) 0.517 48 (59.3%) 33 (40.7%) 0.097 40 (49.4%) 41 (50.6%) 0.120	89 (74.8%) 30 (25.2%) 96 (80.7%) 23 (19.3%) 22 (18.5%) 97 (81.5%) 84 (70.6%) 35 (29.4%) 72 (60.5%) 47 (39.5%)
(5) Use liquid plant oils rather than tropical oils Adherents (n=158, 79.0%) Non-adherents (n=42, 21.0%) p value	10.1±2.3 (10.0) 9.1±2.1 (9.0) 0.008	58 (71.6%) 23 (28.4%) 0.034	100 (84.0%) 19 (16.0%)
(6) Minimize intake of beverages and foods with added sugars Adherents (n=134, 67.0%) Non-adherents (n=66, 33.0%) p value	10.3±2.0 (10.0) 9.1±2.7 (9.0) 0.001	43 (53.1%) 38 (46.9%) 0.001	91 (76.5%) 28 (23.5%)
(7) Choose and prepare foods with little or no salt Adherents (n=168, 84.0%) Non-adherents (n=32, 16.0%) p value	9.9±2.4 (10.0) 9.8±2.1 (10.0) 0.909	70 (86.4%) 11 (13.6%) 0.441	98 (82.4%) 21 (17.6%)
(8) If you do not drink alcohol don't start, if you choose to drink alcohol, limit intake Adherents (n=168, 84.0%) Non-adherents (n=32, 16.0%) p value	9.9±2.2 (10.0) 10.0±2.7 (10.0) 0.925	68 (84.0%) 13 (16.0%) 0.987	100 (84.0%) 19 (16.0%)
(9) Choose minimally processed foods instead of ultra-processed Adherents (n=150, 75.0%) Non-adherents (n=50, 25.0%) p value	10.1±2.2 (10.0) 9.3±2.7 (10.0) 0.133	57 (70.4%) 24 (29.6%) 0.212	93 (78.2%) 26 (21.8%)
(10) Adhere to this guidance wherever food is prepared/consumed Adherents (n=163, 81.5%) Non-adherents (n=37, 18.5%) p value	10.0±2.2 (10.0) 9.1±2.7 (9.0) 0.047	62 (76.5%) 19 (23.5%) 0.136	101 (84.9%) 18 (15.1%)

Main heading of item 4 is "Adequate Healthy plant-Based and Other protein Sources". Accordingly, in the table, the subheadings under this main heading are presented as a, b, c, d1, and d2.

AHA: American Heart Association; TS: Total score; MEDAS: Mediterranean Diet Adherence Scale

Continuous variables, such as MEDAS total scores, were presented as mean±standard deviation (median) and compared using the independent samples t-test for normally distributed data or the Mann-Whitney U test for non-normally distributed data. Categorical variables, such as adherence groups, were expressed as numbers and percentages (n, %) and compared using the chi-square test.

DISCUSSION

Although CVD mortality has declined in recent decades, it remains a major public health concern, especially in areas with poor adherence to preventive behaviors.¹⁴ Modifiable factors like obesity, poor diet, and inactivity cause most CVD cases, with up to 80–90% preventable through lifestyle changes.¹⁵

Awareness of CVD risk factors is a crucial component of disease prevention, influencing dietary choices and lifestyle behaviors.⁹ This study examined the alignment between the Mediterranean diet and 2021 AHA Guidelines, and the link between diet adherence and CVD risk awareness. The findings indicate that individuals with high adherence to the Mediterranean diet have significantly higher levels of CVD risk awareness. Moreover, individuals who follow to the AHA dietary recommendations demonstrate greater adherence to the Mediterranean diet.

The study revealed that individuals with higher adherence to the Mediterranean diet exhibited significantly greater awareness of CVD risk factors, as reflected in their CDRAAQ scores. Subfactors like perceived risk, healthy eating intentions, and behavioral benefits were significantly higher in participants with greater Mediterranean diet adherence. These findings suggest a reciprocal relationship between adherence to the Mediterranean diet and awareness of CVD risk factors. Dietary awareness may encourage healthier food choices and long-term adherence to the Mediterranean diet. These findings are further supported by previous studies indicating that individuals with higher CVD awareness are more likely to adopt healthy lifestyle changes and show improved adherence to healthy dietary approaches.^{16,17} Similarly, a study has shown that a short-term, group-based, multidisciplinary cardiovascular health program delivered by health professionals improves individuals' adherence to the Mediterranean diet. This suggests that educational programs aimed at increasing nutritional awareness may promote adherence to the Mediterranean diet.¹⁸

These findings align with the Knowledge-Attitude-Behavior Model, where behavior change occurs through knowledge, attitude, and action.¹⁹ Understanding how diet affects chronic disease is key to

adopting healthy eating habits.²⁰ In parallel, it has been shown that lack of knowledge is among the barriers to adherence with the Mediterranean diet in adults at high risk for CVD.²¹ However, while knowledge is a fundamental component of behavior change, it is not always sufficient on its own, as other social, psychological, and environmental factors may play a role.¹⁹

The Mediterranean Diet reduces CVD risk factors and improves metabolic health due to its high content of bioactive compounds and antioxidants. It has positive effects on blood pressure, glycemic control, lipid profile, and inflammation, ultimately providing long-term protection against CVD.²² A previous study indicated that individuals with pre-existing medical conditions, such as diabetes, hypertension, and CVD, exhibit better adherence to the Mediterranean diet. A possible explanation for this finding could be that individuals tend to improve their dietary behaviors positively following a CVD diagnosis or heightened perception of risk.²³

Consistent with this perspective, our study suggests that individuals might enhance adherence to the Mediterranean diet after receiving a CVD diagnosis or with increased perceived cardiovascular risk. In line with this, our study found a significant negative correlation between BMI and the perceived benefits and intentions to change sub-dimension of the CDRAAQ ($p < 0.05$), indicating that individuals with greater awareness of the benefits of lifestyle modifications tend to have a lower BMI. Given the average age of CVD patients, having a normal BMI may be especially beneficial. In support of this, another study conducted among university students reported higher median scores on the Cardiovascular Disease Risk Factor Knowledge Level Scale among underweight, overweight, or obese individuals compared to those with normal body weight.²⁴

Differences in results may be partially explained by findings from another study indicating that many adults with high BMI in high-risk populations are aware of their elevated cardiovascular risk due to factors such as obesity and family history of CVD. Nevertheless, it should also be noted that a significant proportion of adults with high BMI remain unaware of their increased cardiovascular risk. Thus, aware-

ness alone may not suffice, emphasizing the need for interventions to translate it into behavior change.²⁵

Furthermore, adherence to the Mediterranean diet was associated with lower DBP ($p<0.05$), but no significant differences were found in SBP between adherence groups ($p>0.05$). These findings suggest that adherence to the Mediterranean diet may facilitate hypertension control in adults with cardiovascular disease. Magriplis et al. observed a 36% reduction in the likelihood of hypertension among individuals with higher adherence to the Mediterranean diet, highlighting its potential role in cardiovascular health.²⁶ A systematic review and metaanalysis found lower SBP in individuals with high Mediterranean diet adherence, though no significant difference in DBP was observed between groups.²⁷ While blood pressure awareness is less studied than obesity as a cardiovascular risk factor, increased awareness has been associated with positive changes in health behaviors, including improved nutrition and physical activity.¹⁷ This study found a correlation between higher awareness of cardiovascular diseases and lower diastolic or SBP ($p<0.05$).

The findings of this study reveal that adherence to specific AHA recommendations is associated with higher MEDAS scores. It was observed that participants who reported maintaining a healthy body weight were more likely to be in the high Mediterranean diet adherence group. Awareness of healthy weight may promote better adherence to the Mediterranean diet. The Mediterranean diet provides a weight loss outcome that is similar to or greater than other diets (such as low-fat and/or low-carbohydrate diets) that promise weight loss but whose long-term health effects are not clearly known.²⁸ The “Prevençión con Dieta Mediterránea (PREDIMED)” and “CORonary Diet Intervention with olive oil and cardiovascular PREvention (CORDIPREV)” studies have confirmed the clear cardiovascular benefits of the Mediterranean diet and have also supported its positive effects on weight loss.^{29,30}

Although no significant association was found between BMI and overall MEDAS scores in this study, the significant negative correlation between the “perceived benefits and intentions for change”

sub-factor of the CDRAAQ and BMI provides an explanation for this. This suggests that awareness of obesity risks may not fully lead to healthy dietary behaviors. Moreover, a previous study evaluating this alignment also reported that the recommendation to maintain a healthy body weight was not scored in the comparison, as it is not included in the Mediterranean diet model.⁷

A detailed examination of dietary habits revealed that adherence to key AHA dietary recommendations was associated with increased adherence to the Mediterranean diet. Specifically, individuals who reported higher and more diverse consumption of fruits and vegetables, preferred fish and seafood, consumed poultry instead of red and processed meats, and favored liquid plant oils over tropical oils demonstrated significantly greater adherence to the Mediterranean diet. These findings align with prior studies on the role of these food groups in cardiovascular health. Furthermore, the results suggest that choosing healthy sources of fats and proteins may also enhance dietary adherence.^{31,32}

It was found that individuals who limited their intake of added sugars had significantly higher MEDAS scores. Although no significant associations were found between MEDAS scores and the limitation of ultra-processed food consumption in this study, previous research showed that higher adherence to the Mediterranean diet was significantly associated with lower consumption of ultra-processed foods. In that study, children in the highest adherence group reported an 8.5% lower energy intake from ultra-processed foods compared to those in the lowest adherence group. Additionally, it was highlighted that 71.6% of the variability in free sugar intake was explained by the variability in ultra-processed food consumption. These findings suggest that ultra-processed food (UPF) consumption plays a significant role in increasing free sugar intake, and reducing UPF consumption could be an effective strategy to enhance adherence to the Mediterranean diet.^{33,34}

Furthermore, no significant associations were found between MEDAS total scores and the consumption of whole grains, dairy product preferences, salt use, alcohol consumption ($p>0.05$). This suggests

that these dietary habits may have a limited impact on adherence to the Mediterranean diet. Consistent with this, previous research indicated that adherence to recommendations regarding salt and alcohol was only partially achieved, while recommendations concerning dairy product consumption were not scored, as dairy is not considered a core component of the Mediterranean diet.⁷

STRENGTHS, LIMITATIONS, AND FUTURE RESEARCH DIRECTIONS

One of the strengths of this study is its comprehensive assessment of dietary adherence using both the Mediterranean diet and AHA guideline frameworks, which provides a more nuanced understanding of dietary behaviors in CVD patients. However, several limitations should be acknowledged. First, the cross-sectional design limits causal interpretations, as it cannot establish whether greater awareness leads to improved adherence or vice versa. Second, the reliance on self-reported dietary intake and risk awareness measures introduces potential recall and social desirability biases. Additionally, since the study sample was drawn from a single clinic, the generalizability of the findings may be limited. Future longitudinal studies incorporating objective biomarkers of dietary adherence and broader, more diverse populations are needed to further elucidate these relationships.

CONCLUSION

These findings add to the evidence linking dietary patterns and cardiovascular health, highlighting the need for targeted interventions to raise risk aware-

ness. Greater knowledge of CVD risk factors and preventive behaviors can promote adherence to cardio-protective diets, particularly the Mediterranean diet. Higher adherence was associated with better risk awareness, lower BMI, and improved blood pressure control.

This study emphasizes the importance of integrating patient education and awareness programs into clinical care. Empowering individuals to make informed dietary choices may improve cardiovascular outcomes. Future research should develop and assess long-term educational strategies to sustain adherence to heart-healthy diets.

Source of Finance

This study was supported way TUBITAK 2209-A-University Students Research Projects Support Program (Project no: 1919B012110788).

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Büşra Başar Gökçen, Dila Özdemir; **Design:** Büşra Başar Gökçen, Dila Özdemir; **Control/Supervision:** Büşra Başar Gökçen; **Data Collection and/or Processing:** Dila Özdemir, Sena Aksu; **Analysis and/or Interpretation:** Büşra Başar Gökçen; **Literature Review:** Büşra Başar Gökçen, Sena Aksu; **Writing the Article:** Büşra Başar Gökçen; **Critical Review:** Büşra Başar Gökçen, Sena Aksu; **References and Fundings:** Dila Özdemir; **Materials:** Dila Özdemir.

REFERENCES

- Ballin M, Neovius M, Ortega FB, Henriksson P, Nordström A, Berglind D, et al. Genetic and environmental factors and cardiovascular disease risk in adolescents. *JAMA Netw Open*. 2023;6(11):e2343947. [Crossref] [PubMed] [PMC]
- Ojeda-Granados C, Campisi E, Barchitta M, Agodi A. Genetic, lifestyle and metabolic factors contributing to cardiovascular disease in the Italian population: a literature review. *Front Nutr*. 2024;11:1379785. [Crossref] [PubMed] [PMC]
- Pietri P. Dietary guidance for cardiovascular health: consensus and controversies. *nutrients*. 2023;15(19):4295. [PubMed] [PMC]
- Fegers-Wustrow I, Gianos E, Halle M, Yang E. Comparison of American and European Guidelines for primary prevention of cardiovascular disease: JACC guideline comparison. *J Am Coll Cardiol*. 2022;79(13):1304-13. [PubMed]
- Visseren FLJ, Mach F, Smulders YM, Carballo D, Koskinas KC, Böck M, et al; ESC National Cardiac Societies; ESC Scientific Document Group. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J*. 2021;42(34):3227-337. Erratum in: *Eur Heart J*. 2022;43(42):4468. [Crossref] [PubMed]
- Lichtenstein AH, Appel LJ, Vadiveloo M, Hu FB, Kris-Etherton PM, Rebholz CM, et al. 2021 Dietary Guidance to improve cardiovascular health: a scientific statement from the American Heart Association. *Circulation*. 2021;144(23):e472-e487. [Crossref] [PubMed]
- Gardner CD, Vadiveloo MK, Petersen KS, Anderson CAM, Springfield S, Van Horn L, et al; American Heart Association Council on Lifestyle and Cardiometabolic Health. Popular dietary patterns: alignment with American Heart Association 2021 dietary guidance: a scientific statement from the American Heart Association. *Circulation*. 2023;147(22):1715-30. [Crossref] [PubMed]
- Jiva ES. Atherosclerosis-preventing CVD by increasing awareness regarding risk factors. *Journal of Clinical Lipidology*. 2022;16(1):e32-e4. [Crossref] [PubMed]
- Qasem Surrafi AM, Mohammedsaeed W, Shikieri ABE. Cardiovascular risk awareness and calculated 10-year risk among female employees at Taibah University 2019. *Front Public Health*. 2021;9:658243. [PubMed] [PMC]
- Woringer M, Nielsen JJ, Zibarras L, Evason J, Kassianos AP, Harris M, et al. Development of a questionnaire to evaluate patients' awareness of cardiovascular disease risk in England's National Health Service Health Check preventive cardiovascular programme. *BMJ Open*. 2017;7(9):e014413. [Crossref] [PubMed] [PMC]
- Vural Doğru B, Utlı H, Karaman E. Cardiovascular Disease Risk Awareness Assessment Questionnaire: psychometric properties of the Turkish Version. *Türk J Card Nur*. 2021;12(27):18-25. [Crossref]
- Martínez-González MA, García-Arellano A, Toledo E, Salas-Salvadó J, Buil-Cosiales P, Corella D, et al; PREDIMED study investigators. a 14-item Mediterranean diet assessment tool and obesity indexes among high-risk subjects: the PREDIMED trial. *PLoS One*. 2012;7(8):e43134. [Crossref] [PubMed] [PMC]
- Özkan Pehlivanoglu EF, Balcioglu H, Ünlüoglu İ. Akdeniz Diyeti Bağlılık Ölçeği'nin Türkçe'ye uyarlanması geçerlilik ve güvenilirliği [Turkish validation and reliability of Mediterranean Diet Adherence Screener]. *Osmangazi Tıp Dergisi*. 2020;42(2):160-4. [Link]
- Sozmen K, Ergor G, Sakarya S, Dinc Horasan G, Sahan C, Ekinci B, et al. Evaluation of blood pressure status and mortality in Turkey: findings from chronic diseases and risk factors cohort study. *Medicina (Kaunas)*. 2023;59(8):1366. [PubMed] [PMC]
- Güneş FE, Bekiroglu N, Imeryuz N, Agirbasli M. Awareness of cardiovascular risk factors among university students in Turkey. *Prim Health Care Res Dev*. 2019;20:e127. Erratum in: *Prim Health Care Res Dev*. 2019;20:e148. [Crossref] [PubMed] [PMC]
- Sulaiman HA, Andsoy II. Health behaviors, knowledge, and attitudes toward cardiovascular disease risk factors in young Iraqi adults: a sample from Erbil, Iraq. *Cardiovasc Prev Pharmacother*. 2024;6(3):92-101. [Crossref]
- Jacobs JC, Burke S, Rouse M, Sarma S, Zaric G. Cardiovascular disease risk awareness and its association with preventive health behaviors: evidence from a sample of Canadian workplaces. *J Occup Environ Med*. 2016;58(5):459-65. [PubMed]
- Mosher LT, Bizerra C, Davies K, Seabrook JA, Keathley J. Evaluation of a 4-week interdisciplinary primary care cardiovascular health programme: impact on knowledge, Mediterranean diet adherence and biomarkers. *BMJ Nutr Prev Health*. 2024;7(1):95-102. [Crossref] [PubMed] [PMC]
- Turner A, LaMonica HM, Moroney C, O'Leary F, Naismith SL, Flood VM. Knowledge, attitudes, and behaviours concerning the Mediterranean diet among older adults in Australia. *J Community Health*. 2023;48(6):951-62. [Crossref] [PubMed] [PMC]
- Peltzer K, Pengpid S. Prevalence, risk awareness and health beliefs of behavioural risk factors for cardiovascular disease among university students in nine ASEAN countries. *BMC Public Health*. 2018;18(1):237. [PubMed] [PMC]
- Moore SE, McEvoy CT, Prior L, Lawton J, Patterson CC, Kee F, et al. Barriers to adopting a Mediterranean diet in Northern European adults at high risk of developing cardiovascular disease. *J Hum Nutr Diet*. 2018;31(4):451-62. [PubMed]
- Richardson LA, Izuora K, Basu A. Mediterranean diet and its association with cardiovascular disease risk factors: a scoping review. *Int J Environ Res Public Health*. 2022;19(19):12762. [Crossref] [PubMed] [PMC]
- Zeenny RM, Haddad C, Hajj A, Zeidan RK, Salameh P, Ferrières J. Adherence to the Mediterranean diet and cardiovascular risk factors among the Lebanese population: a nationwide cross-sectional post hoc study. *Nutrients*. 2024;16(15):2426. [PubMed] [PMC]
- Kes D, Sanlitürk D, Polat U. University students' knowledge levels about cardiovascular risk factors and assessment of their health behaviours in Turkey. *Int J Caring Sci*. 2018;11(2):1269-81. [Link]
- Petrovic J, Radomirovic M, Petrovic Djordjevic I, Petrovic A, Markovic M, Todorovic J, et al. Awareness of personal cardiovascular risk among the adults with an elevated body mass index in a very high-risk country. *Eur Heart J*. 2024;45(Supplement_1):ehae666. 2896. [Crossref]
- Magriplis E, Panagiotakos D, Kyrou I, Tsioufis C, Mitsopoulou AV, Karageorgou D, et al. Presence of hypertension is reduced by Mediterranean diet adherence in all individuals with a more pronounced effect in the obese: the Hellenic National Nutrition and Health Survey (HNNHS). *Nutrients*. 2020;12(3):853. [Crossref] [PubMed] [PMC]
- Bakaloudi DR, Chrysoula L, Leonida I, Kotzakioulafi E, Theodoridis X, Chourdakis M. Impact of the level of adherence to the Mediterranean diet on blood pressure: a systematic review and meta-analysis of observational studies. *Clin Nutr*. 2021;40(12):5771-80. [Crossref] [PubMed]
- Dominguez LJ, Veronese N, Di Bella G, Cusumano C, Parisi A, Tagliaferri F, et al. Mediterranean diet in the management and prevention of obesity. *Exp Gerontol*. 2023;174:112121. [Crossref] [PubMed]
- Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, et al; PREDIMED study investigators. primary prevention of cardiovascular disease with a Mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med*. 2018;378(25):e34. [Crossref] [PubMed]
- Delgado-Lista J, Alcalá-Díaz JF, Torres-Peña JD, Quintana-Navarro GM, Fuentes F, García-Ríos A, et al; CORDIOPREV Investigators. Long-term secondary prevention of cardiovascular disease with a Mediterranean diet and a low-fat diet (CORDIOPREV): a randomised controlled trial. *Lancet*. 2022;399(10338):1876-85. [PubMed]

31. Godoy-Izquierdo D, Ogallar A, Lara R, Rodríguez-Tadeo A, Arbinaga F. Association of a Mediterranean diet and fruit and vegetable consumption with subjective well-being among adults with overweight and obesity. *Nutrients*. 2021;13(4):1342. [[PubMed](#)] [[PMC](#)]
32. Marinac Pupavac S, Kenđel Jovanović G, Linšak Ž, Glad M, Traven L, Pavičić Žeželj S. The influence on fish and seafood consumption, and the attitudes and reasons for its consumption in the Croatian population. *Front Sustain Food Syst*. 2022;6:945186. [[Crossref](#)]
33. da Rocha BRS, Rico-Campà A, Romanos-Nanclares A, Ciriza E, Barbosa KBF, Martínez-González MÁ, et al. Adherence to Mediterranean diet is inversely associated with the consumption of ultra-processed foods among Spanish children: the SENDO project. *Public Health Nutr*. 2021;24(11):3294-303. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
34. Dinu M, Tristan Asensi M, Pagliai G, Lotti S, Martini D, Colombini B, et al. Consumption of ultra-processed foods is inversely associated with adherence to the Mediterranean diet: a cross-sectional study. *Nutrients*. 2022;14(10):2073. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]