

ORIGINAL RESEARCH ORJİNAL ARAŞTIRMA

DOI: 10.5336/nurses.2024-105012

Healthy Lifestyle Behaviors of Primary School-Aged Children and Affecting Predictors: A Cross-Sectional Study

İlkokul Çağındaki Çocukların Sağlıklı Yaşam Biçimi Davranışları ve Etkileyen Yordayıcılar: Kesitsel Çalışma

^aEda KILINÇ İŞLEYEN^a, ^bEdanur ÖZKAYA BOZKURT^b, ^cAsiye KARTAL^b

^aUşak University Faculty of Health Science, Department of Nursing, Department of Public Health Nursing, Uşak, Türkiye

^bPamukkale University Faculty of Health Science, Department of Nursing, Department of Public Health Nursing, Denizli, Türkiye

ABSTRACT Objective: To develop effective interventions to improve children's health outcomes, it is of great importance to determine the health-related lifestyles of primary school-aged children and the factors that influence these lifestyles. The aim of this research was to identify the elements that influence primary school children's healthy lifestyle behaviors. **Material and Methods:** This is a cross-sectional study. The research 1,240 primary school-aged children who met the inclusion criteria formed the sample of the study. Descriptive characteristics form and Healthy Lifestyle Behaviors in Children Scale (HLBCS) were used. Data collection was carried out in the school-aged children's own classrooms using face-to-face data collection technique. Analysis of variance, t-test, and multiple linear regression analysis was used. **Results:** Female school-aged children, school-aged children whose parents are living, school-aged children whose mother has a university education, whose father has a high school or university education, who has a good economic situation and who does not have an overweight family member, have significantly higher mean HLBCS scores. **Conclusion:** It was shown that primary school-aged children had a good level of lifestyle behaviors. The most important predictors affecting lifestyle behaviors were gender, mother's education level, income level and class level. To improve school-aged children's lifestyle behaviors, it is important to provide education on physical activity, nutrition, hygiene and sleep by the school nurse.

Keywords: Healthy lifestyle; health behavior; child health; school health; school health nursing

ÖZET Amaç: Çocukların sağlık sonuçlarını iyileştirmeye yönelik etkili müdahaleler geliştirmek için, ilkökul çağındaki çocukların sağlıklı yaşam biçimlerini ve bu yaşam biçimlerini etkileyen faktörleri belirlemek büyük önem taşımaktadır. Bu araştırmanın amacı, ilkökul çağındaki çocukların sağlıklı yaşam biçimi davranışlarını etkileyen faktörlerin belirlenmesidir. **Gereç ve Yöntemler:** Bu araştırma kesitseldir. Araştırmanın örneklemini 1.240 ilkökul çağındaki çocuk oluşturmaktadır. Tanımlayıcı Özellikler Formu ve Çocuklarda Sağlıklı Yaşam Biçimi Davranışları Ölçeği kullanılmıştır. Veri toplama, okul çağı çocuklarının kendi sınıflarında yüz yüze veri toplama tekniği kullanılarak gerçekleştirilmiştir. Varyans analizi, t-testi ve çoklu doğrusal regresyon analizi kullanılmıştır. **Bulgular:** Kız öğrencilerin, anne ve babası hayatta olan öğrencilerin, annesi üniversite eğitimi almış olan öğrencilerin, babası lise veya üniversite eğitimi almış olan öğrencilerin, ekonomik durumu iyi olan öğrencilerin ve ailesinde aşırı kilolu birey olmayan öğrencilerin Çocuklarda Sağlıklı Yaşam Biçimi Davranışları Ölçeği puan ortalamaları anlamlı düzeyde daha yüksektir. **Sonuç:** İlkokul çağı çocuklarının iyi düzeyde yaşam biçimi davranışlarına sahip olduğu gösterilmiştir. Yaşam biçimi davranışlarını etkileyen en önemli yordayıcılar cinsiyet, anne eğitim düzeyi, gelir düzeyi ve sınıf düzeyidir. Çocukların yaşam biçimi davranışlarını geliştirmek için okul hemşiresi tarafından fiziksel aktivite, beslenme, hijyen ve uyku konularında eğitim verilmesi önemlidir.

Anahtar Kelimeler: Sağlıklı yaşam biçimi; sağlık davranışı; çocuk sağlığı; okul sağlığı; okul sağlığı hemşireliği

Healthy lifestyle behaviors are characterized as all behaviors describe that have an impact on a person's health and can be controlled against factors that have a significant impact on their health.¹ In other

words, it refers to the entirety of the actions and beliefs a person adopts to maintain their health and ward off illness.² When trying to enhance health, one public health policy that should be taken into considera-

TO CITE THIS ARTICLE:

Kılınç İşleyen E, Özkaya Bozkurt E, Kartal A. Healthy lifestyle behaviors of primary school-aged children and affecting predictors: A cross-sectional study. Türkiye Klinikleri J Nurs Sci. 2025;17(2):436-46.

Correspondence: Edanur ÖZKAYA BOZKURT

Pamukkale University Faculty of Health Science, Department of Nursing, Department of Public Health Nursing, Denizli, Türkiye

E-mail: edaozkaya19@gmail.com

Peer review under responsibility of Türkiye Klinikleri Journal of Nursing Sciences.

Received: 31 Jul 2024

Received in revised form: 17 Oct 2024

Accepted: 14 Nov 2024

Available online: 17 Mar 2025

2146-8893 / Copyright © 2025 by Türkiye Klinikleri. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



tion is the promotion of a healthy lifestyle.³ Healthy lifestyle behaviors are summarized as having adequate and balanced nutritional habits, managing stress, doing regular physical activity, engaging in spiritual activities, keeping interpersonal relationships alive and taking responsibility in these matters.⁴ The acquisition of good lifestyle habits from childhood is crucial for the long-term maintenance of those behaviors.¹

It is especially important to promote healthy behaviors early on since childhood habits carry over into adulthood. Lifestyle choices during this period can result in a number of health issues in adulthood. Increased harmful habits including childhood smoking, physical inactivity, and malnutrition are contributing to the rise in lifestyle-related health issues like diabetes, heart disease, and obesity.⁵ Children who are still going through a period of physical, mental, and emotional development are therefore especially susceptible to these consequences.⁶ This highlights even more how critical it is to establish healthy habits early in infancy to promote appropriate growth and development.

Instilling positive health behaviors in children at an early age is extremely important for public health as it ensures health and well-being in adulthood and improves physical health and academic performance.⁷ Adopting healthy habits can also help children feel better about themselves and have less mental health issues.⁸ There is great potential in using school environments to improve health habits and positively affect academic success.⁹ The child has to adopt a healthy lifestyle in all areas, such as diet, exercise, behavioral development, and entertainment, in order to grow up to be a healthy adult.¹⁰

In a systematic review examining lifestyle behaviors in children, it was determined that unhealthy nutrition, high levels of inactivity and sedentary lifestyle were observed in children.¹¹ In a research study with comprehensive data collected by World Health Organization from 25 countries, information on physical activity behavior, screen time and sleep time of 150,651 children was analyzed. The data of this study showed that 79.4% of children actively play digital games for more than 1 hour every day,

53.9% are not members of a sports or dance club, and 39.8% spend more than 2 hours in front of the screen a day.¹² In their research, Lekše et al. found an important theme that children need education about health promotion and health behaviors.¹³ These comprehensive data show that primary school-aged children's healthy lifestyle behaviors are below the ideal level and that they require more instruction in this area.

While reviewing the literature, it became evident that many studies on healthy lifestyle behaviors focused on adolescents, but studies on the subject in primary school-aged children were insufficient.^{1,3,14} There is no research in the literature that examines all aspects of healthy lifestyle behaviors in primary school-aged children.¹⁵ But early implementation of treatments aiming at fostering better lives may be more effective in preventing harmful choices from being ingrained in an individual's lifestyle.¹⁶ To improve children's health outcomes, it is essential to identify the health-related lifestyles of elementary school-aged children and the factors that impact these lifestyles. The school health nurse plays a very important role in developing healthy lifestyle behaviors in primary school-aged children's. This role is critical for children to acquire healthy habits and adopt a long-term healthy lifestyle.¹⁷

Thus, the purpose of this study was to identify the characteristics that influence primary school-aged children's adoption of healthy lifestyles. It is expected that the study results will guide future studies on the subject and contribute to the literature.

Research Questions

What are the healthy lifestyle behavior scores of primary school-aged children's?

Is there a difference in healthy lifestyle behavior scores of primary school-aged children's according to their sociodemographic characteristics?

What are the factors that affect primary school-aged children's healthy lifestyle behaviors?

MATERIAL AND METHODS

DESIGN

This research is cross-sectional.

STUDY SETTING AND SAMPLE

This study was conducted between January and May 2023 in the city center of a province located in the west of Türkiye. The population of the research consisted of 3rd and 4th grade school-aged children's studying in 7 primary schools located in the city center affiliated with the Provincial Directorate of National Education. These schools were determined by simple random sampling method among 47 primary schools in the city center. Criteria for inclusion in the sample were being a 3rd or 4th grade primary school-aged children and parental consent. Exclusion criteria are having a mental or physical disability, being a foreign school-aged children and not knowing Turkish. In the sample calculation of this study, the average score of healthy lifestyle behaviors of primary school-aged children's was taken as reference in the study conducted by Öcal and Önsüz.¹⁸ The average score of school-aged children's healthy lifestyle behaviors is 141.39 ± 16.21 . It was assumed that the average score of healthy lifestyle behaviors of the sample in our study was 139.80. Accordingly, effect size=0.098. In the calculation, the sample size was calculated as at least 1,127 by taking the type 1 error margin (α) as 0.05 and the power ($1-\beta$) as 0.95. Considering the possibility of data loss in the research, 1,240 primary school-aged children's who met the inclusion criteria formed the sample of the study.

INSTRUMENTS

Descriptive characteristics form and Healthy Lifestyle Behaviors Scale in Children were utilized as data research collection instruments in the study.

DESCRIPTIVE CHARACTERISTICS FORM

The form with the descriptive characteristics was created considering the literature.^{1,3,18} The form includes questions for school-aged children's including their age, grade, gender, and number of siblings. In addition, there are questions regarding the demographic information of the parents, including the living situation, cohabitation status, education level and profession of the mother and father. In addition, the family was asked about their family type, economic status and whether there were any overweight individuals.

HEALTHY LIFESTYLE BEHAVIORS IN CHILDREN SCALE

This scale was developed by Öcal and Önsüz to measure the healthy lifestyle behaviors of primary school-aged children's in Türkiye.¹⁸ Healthy Lifestyle Behaviors in Children Scale (HLBCS) consists of 53 items and 7 sub-dimensions. These sub-dimensions and Cronbach alpha values are as follows: nutrition sub-dimension, physical activity sub-dimension, hygiene sub-dimension, sleep sub-dimension, environmental cleanliness dimension, health responsibility dimension and addiction dimension. Goodness of fit indexes of the scale are $\chi^2/df=2.55$, Root Mean Square Error of Approximation=0.055, Comparative Fit Index=0.802, Goodness of Fit Index=0.790. The options of the items are scored as "I agree 3, I am undecided 2 and I disagree 1". In each question, children are asked to mark only one of the 3 options that is most appropriate for them. The minimum score that can be obtained from the scale is 53, while the maximum score is 159. As the score obtained from the scale increases, the level of school-aged children's 'healthy lifestyle behaviors also increases. In this study, the Cronbach's Alpha value of the scale was found to be 0.89.

PROCEDURE

School-aged children's who achieved the criteria for participation were given information on the aim and details of the research. Before the data collection process, school administrators and teachers of the relevant classes were contacted, and the school-aged children's parents were contacted, and their informed consent was obtained. Data collection was carried out in the school-aged children's own classrooms using face-to-face data collection technique. Since the survey form was based on self-report, surveys were distributed to school-aged children's, and they were given until the end of the course (40 minutes) to fill them out. During this period, the researchers were in the relevant classroom and answered the school-aged children's questions.

STATISTICAL ANALYSIS

The data were evaluated in the IBM SPSS v29.0 (IBM Corp., Armonk, NY, USA) statistical program.

Count, proportion, average, standard deviation, and lowest-and-maximum values were employed in the descriptive variable analysis. Normal distribution was tested with Skewness=1.02 and Kurtosis=1.49 values and it was determined that the data was normally distributed.¹⁹ Independent sample t-test was used to determine whether there was a difference between the healthy lifestyle behaviors scale mean score between 2 groups in terms of sociodemographic characteristics, and one-way analysis of variance was used to determine the difference in mean score between 3 or more groups. Variables with variations across groups were included in multiple linear regression analysis to identify the predictors influencing healthy lifestyle behaviors. A p value of less than 0.05 was accepted as statistically significant in every analysis.

ETHICAL PRINCIPLES AND APPROVAL

This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Non-Interventional Clinical Research Ethics Committee (Date: December 07, 2022/No: E-60116787-020-301347). Explanations were made to the students to be included in the research and their parents about the purpose of the research and the content of the forms, and informed parental consent was obtained for their participation. Written permissions were obtained from the ministries and institutions to which the data will be collected. Permission was also obtained from the researchers developing the scales used via e-mail.

RESULTS

PARTICIPANT CHARACTERISTICS

Primary school-aged children's of 50.2% are boys and 49.8% are girls. The school-aged children's of 51.0% are in the 3rd grade and 49.0% are in the 4th grade. When the body mass index (BMI) of the school-aged children's was calculated, it was determined that 72.1% were normal weight, 10.6% were overweight, 8.7% were underweight and 8.6% were obese (Table 1). In the BMI classification, the values determined by Neyzi et al. for Turkish children were taken as reference.²⁰

TABLE 1: Demographic characteristics of the students

| | n | % |
|--|-------|-------|
| Gender | | |
| Female | 617 | 49.8 |
| Male | 623 | 50.2 |
| Age | | |
| 8 | 220 | 17.7 |
| 9 | 550 | 44.4 |
| 10 | 470 | 37.9 |
| Grade | | |
| 3 rd | 633 | 51.0 |
| 4 th | 607 | 49.0 |
| Number of siblings | | |
| None | 199 | 16.0 |
| 1 | 495 | 39.9 |
| 2 | 319 | 25.7 |
| 3 and above | 227 | 18.3 |
| Living situation of mother and father | | |
| Both are alive | 1,214 | 97.9 |
| Only the mother lives | 20 | 1.6 |
| Only the father lives | 5 | 0.4 |
| Both passed away | 1 | 0.1 |
| Living situation of mother and father | | |
| Living together | 1,112 | 89.7 |
| Living apart-death | 128 | 8.6 |
| Mother's educational status | | |
| Illiterate | 93 | 7.5 |
| Primary school | 209 | 16.9 |
| Middle school | 229 | 18.5 |
| High school | 290 | 23.4 |
| University | 419 | 33.8 |
| Father's educational status | | |
| Illiterate | 74 | 6.0 |
| Primary school | 156 | 12.6 |
| Middle school | 235 | 19.0 |
| High school | 336 | 27.1 |
| University | 439 | 35.4 |
| Mother's profession | | |
| Housewife | 605 | 48.8 |
| Officer | 261 | 21.0 |
| Employee | 333 | 26.9 |
| Other occupations and death | 41 | 3.3 |
| Father's profession | | |
| Not working | 69 | 5.6 |
| Officer | 349 | 28.1 |
| Employee | 748 | 60.3 |
| Other occupations and death | 74 | 6.0 |
| Economic level of the family | | |
| Income is less than expenses | 781 | 63.0 |
| Income is equal to expenses | 430 | 34.7 |
| Income is more than expenses | 29 | 2.30 |
| Family type | | |
| Nuclear family | 913 | 73.6 |
| Extended family | 327 | 26.4 |
| BMI Percentile Value Classification (kg/m ²) | | |
| <5 | 108 | 8.70 |
| 5-15 | 208 | 16.80 |
| 15-25 | 134 | 10.80 |
| 25-50 | 258 | 20.80 |
| 50-75 | 201 | 16.20 |
| 75-85 | 93 | 7.50 |
| 85-95 | 131 | 10.6 |
| 95> | 107 | 8.60 |
| Overweight level according to BMI | | |
| <5 (Low weight) | 108 | 8.70 |
| 5-85 (Normal weight) | 894 | 72.10 |
| 85-95 (Overweight) | 131 | 10.60 |
| 95> (Obese) | 107 | 8.60 |
| Is there an overweight member in the family? | | |
| Yes | 550 | 44.4 |
| No | 690 | 55.6 |

BMI: Body mass index

THE MEAN SCORES ON THE HEALTHY LIFESTYLE BEHAVIORS SCALE

The mean score of the primary school-aged children's on the HLBCS was 140.96 ± 12.23 . The mean scores of the HLBCS subscales are between 4.93 and 48.26 (Table 2).

THE COMPARISON OF THE MEAN SCORES ON THE HEALTHY LIFESTYLE BEHAVIOURS SCALE BETWEEN SOCIODEMOGRAPHIC CHARACTERISTICS

Table 3 illustrates the statistically significant variations that were discovered between gender, health status of the parents, education level of the mother and father, economic status, and the status of being an overweight member in the family and the average score of healthy lifestyle behaviors ($p < 0.05$). Female school-aged children, school-aged children's whose parents are living, school-aged children's whose mother has a university education, whose father has a high school or university education, who has a good economic situation and who does not have an overweight family member, have significantly higher mean HLBCS scores ($p < 0.05$) (Table 3).

PREDICTORS AFFECTING HEALTHY LIFESTYLE BEHAVIORS IN PRIMARY SCHOOL-AGED CHILDREN'S

Predictors affecting healthy lifestyle behaviors in primary school-aged children's are gender, grade, maternal education level and economic status. Being a female gender ($\beta = 0.155$), being a 3rd grade school-aged children's ($\beta = 0.123$), having a mother's education level of university or higher ($\beta = 0.143$), and

having a good economic level ($\beta = 0.135$) positively affect healthy lifestyle behaviors ($p < 0.05$) (Table 4).

DISCUSSION

Primary school age is a very important period for the development of healthy behaviors that will affect one's health and well-being in the later stages of life. For this reason, this study primarily examined the level of healthy lifestyle behaviors of primary school-aged children's and the factors affecting these behaviors.

The mean score of the primary school-aged children's included in this study was 140.96 ± 12.23 . Accordingly, our result shows that the school-aged children's participating in the study have high healthy lifestyle behaviors score. In the literature, primary school-aged children's and adolescents have similar problems with risky behaviors such as malnutrition habits, inadequate physical activity, unhealthy leisure time activities, alcohol and tobacco use that can lead to health problems and have low healthy lifestyle behaviors.^{6,12,13,15,21-23} Contrary to the literature, our research finding showed that the lifestyle behaviors of the primary school-aged children's included in the sample were at a positive level. While this result is considered positive, it suggests that the reason for this may be related to family, environment, school environment and role modeling.

The fact that the majority of the parents of the children included in the sample were university graduates may have had a positive effect on the result. Parents' lifestyle behaviors are one of the main factors affecting children's behaviors. As stated by Lekše et al. the most basic need of primary school children is education on health promotion and healthy behaviors.¹³ Schools' healthy nutrition policies, programs encouraging physical activity and health-oriented activities can support school-aged children's healthy lifestyle behaviors. The fact that the schools where this study was conducted were located in the city center and had facilities and that planned trainings were given by teachers and nursing school-aged children's on these issues may have positively affected the result.

In this study, the predictors affecting lifestyle behaviors were also examined. According to the results

TABLE 2: Mean scores on the healthy lifestyle behaviors scale

| Sub-Dimensions | \bar{X} | SD | Minimum | Maximum |
|-----------------------|-----------|-------|---------|---------|
| Nutrition | 48.26 | 5.61 | 19.00 | 57.00 |
| Physical activity | 10.06 | 1.84 | 4.00 | 12.00 |
| Hygiene | 22.71 | 1.78 | 8.00 | 24.00 |
| Sleep | 4.93 | 1.04 | 2.00 | 6.00 |
| Environment | 16.81 | 1.71 | 6.00 | 18.00 |
| Health responsibility | 24.76 | 2.65 | 9.00 | 27.00 |
| Dependence | 13.39 | 1.82 | 5.00 | 15.00 |
| HLBCS total score | 140.96 | 12.23 | 53.00 | 159.00 |

SD: Standard deviation; HLBCS: Healthy Lifestyle Behaviors in Children Scale

TABLE 3: Relationship between HLBCS and sociodemographic characteristics

| | Nutrition | Physical activity | Hygiene | Sleep | Environmental cleanliness | Health responsibility | Addiction | Total |
|---------------------------------------|----------------------|----------------------|----------------------|----------------------|---------------------------|-----------------------|----------------------|------------------------|
| Gender | | | | | | | | |
| Female | 48.72±5.60 | 9.76±1.89 | 23.07±1.45 | 5.05±1.00 | 17.08±1.47 | 25.03±2.29 | 13.64±1.69 | 142.38±11.50 |
| Male | 47.81±5.58 | 10.36±1.74 | 22.36±2.00 | 4.82±1.08 | 16.54±1.87 | 24.49±2.95 | 13.13±1.91 | 139.55±12.77 |
| t-test/p value | 2.880/p=0.004 | -5.805/p=0.000 | 7.121/p=0.000 | 3.799/p=0.000 | 5.652/p=0.000 | 3.573/p=0.000 | 4.935/p=0.000 | 4.106 p=0.000 |
| Age | | | | | | | | |
| 8 | 49.42±5.20 | 10.09±1.88 | 22.75±1.78 | 5.08±0.95 | 16.95±1.49 | 24.83±2.60 | 13.57±1.84 | 142.70±11.05 |
| 9 | 48.31±5.83 | 10.05±1.84 | 22.74±1.84 | 4.97±1.04 | 16.86±1.64 | 24.88±2.61 | 13.50±1.75 | 141.33±12.62 |
| 10 | 47.67±5.44 | 10.07±1.83 | 22.67±1.72 | 4.82±1.08 | 16.68±1.86 | 24.58±2.73 | 13.18±1.88 | 139.70±12.18 |
| Anova/p value | 7.398 p=0.001 | 0.029/p=0.971 | 0.190/p=0.827 | 5.161/p=0.006 | 2.282/p=0.102 | 1.678/p=0.187 | 5.212 p=0.006 | 5.014 p=0.007 |
| | 8 vs 10<0.05 | | | 8 vs 10<0.05 | | | 8 vs 10<0.05 | 8 vs 10<0.05 |
| | 8 vs 9<0.05 | | | | | | 9 vs 10<0.05 | |
| Grade | | | | | | | | |
| 3 rd | 48.95±5.56 | 10.26±1.77 | 22.79±1.77 | 5.08±0.97 | 16.97±1.49 | 24.96±2.51 | 13.56±1.76 | 142.60±11.47 |
| 4 th | 47.55±5.57 | 9.86±1.89 | 22.63±1.79 | 4.78±1.10 | 16.63±1.89 | 24.55±2.78 | 13.21±1.87 | 139.24±12.76 |
| t-test/p value | 4.410/p=0.000 | 3.857/p=0.000 | 1.530/p=0.126 | 5.170/p=0.000 | 3.484/p=0.000 | 2.738/p=0.006 | 3.401/p=0.001 | 4.872 p=0.000 |
| Family type | | | | | | | | |
| Nuclear family | 48.39±5.53 | 10.04±1.85 | 22.76±1.73 | 4.97±1.04 | 16.87±1.59 | 24.85±2.58 | 13.43±1.82 | 141.34±11.99 |
| Extended family | 47.92±5.80 | 10.13±1.80 | 22.60±1.94 | 4.83±1.05 | 16.62±1.99 | 24.50±2.85 | 13.25±1.83 | 139.88±12.84 |
| t test/p value | 1.293/p=0.196 | -0.719/p=0.473 | 1.368/p=0.172 | 2.026/p=0.043 | 2.093/p=0.037 | 1.932/p=0.054 | 1.523/p=0.128 | 1.851/p=0.064 |
| Number of siblings | | | | | | | | |
| None | 47.74±5.80 | 9.73±2.14 | 22.64±1.90 | 4.90±1.07 | 16.63±1.99 | 24.80±2.77 | 13.24±1.97 | 139.71±13.87 |
| 1 | 48.79±5.72 | 10.16±1.84 | 22.74±1.81 | 5.00±1.03 | 16.88±1.65 | 24.78±2.57 | 13.45±1.83 | 141.84±12.22 |
| 2 | 48.36±5.45 | 10.11±1.71 | 22.73±1.67 | 4.91±1.06 | 16.84±1.63 | 24.79±2.55 | 13.50±1.69 | 141.29±11.54 |
| 3 and above | 47.43±5.30 | 10.08±1.71 | 22.70±1.78 | 4.85±1.04 | 16.76±1.65 | 24.61±2.89 | 13.22±1.84 | 139.66±11.52 |
| Anova/p value | 3.779/p=0.010 | 2.727/p=0.043 | 0.173/p=0.915 | 1.328/p=0.264 | 1.048/p=0.370 | 0.299/p=0.826 | 1.757/p=0.154 | 2.479/p=0.060 |
| | 1 vs 3<0.05 | 0 vs 1<0.05 | | | | | | |
| Living situation of mother and father | | | | | | | | |
| Both are alive | 48.26±5.59 | 10.07±1.83 | 22.72±1.76 | 4.93±1.04 | 16.83±1.67 | 24.78±2.64 | 13.40±1.82 | 141.02±12.09 |
| Only the mother lives | 49.50±4.33 | 9.80±2.14 | 23.05±1.98 | 5.30±0.86 | 16.50±1.87 | 24.40±2.70 | 13.00±1.80 | 141.55±11.45 |
| Only the father lives | 44.40±12.34 | 9.00±3.00 | 20.40±4.03 | 4.60±1.51 | 14.60±4.92 | 23.20±3.89 | 12.00±2.34 | 128.20±30.55 |
| Both passed away | 42.00±0.00 | 7.00±0.00 | 18.00±0.00 | 3.00±0.00 | 12.00±0.00 | 18.00±0.00 | 11.00±0.00 | 111.00±0.00 |
| Anova/p value | 1.531/p=0.205 | 1.639/p=0.179 | 5.410/p=0.001 | 2.109/p=0.097 | 5.759/p=0.001 | 2.888/p=0.035 | 1.874/p=0.132 | 3.867 p=0.009 |
| Living situation of mother and father | | | | | | | | |
| Living together | 48.26±5.58 | 10.08±1.82 | 22.71±1.76 | 4.93±1.04 | 16.83±1.70 | 24.79±2.62 | 13.42±1.80 | 141.03±12.16 |
| Living apart-death | 48.29±5.98 | 9.96±2.07 | 22.82±1.90 | 4.99±1.07 | 16.71±1.73 | 24.65±2.93 | 13.14±1.98 | 140.59±12.72 |
| Anova/p value | 0.017/p=0.983 | 0.507/p=0.602 | 0.378/p=0.686 | 0.186/p=0.831 | 1.626/p=0.197 | 1.647/p=0.193 | 1.359/p=0.257 | 0.407/p=0.666 |
| Mother's educational status | | | | | | | | |
| Illiterate | 47.68±5.92 | 9.68±1.96 | 22.40±2.21 | 4.66±1.11 | 16.67±1.83 | 24.09±2.88 | 13.33±1.79 | 138.55±13.17 |
| Primary school | 47.43±5.51 | 9.99±1.92 | 22.59±1.92 | 4.91±1.04 | 16.45±1.91 | 24.29±2.88 | 13.15±1.98 | 138.84±12.44 |
| Middle school | 48.11±5.49 | 9.92±1.90 | 22.88±1.54 | 4.96±1.03 | 16.92±1.58 | 24.86±2.49 | 13.52±1.68 | 141.19±11.17 |
| High school | 48.17±5.69 | 10.14±1.83 | 22.75±1.68 | 4.91±1.06 | 16.81±1.74 | 24.96±2.48 | 13.34±1.83 | 141.11±12.10 |
| University | 48.96±5.53 | 10.21±1.74 | 22.73±1.80 | 5.00±1.02 | 16.95±1.59 | 24.94±2.64 | 13.48±1.82 | 142.31±12.39 |
| Anova/p value | 3.084/p=0.015 | 2.236/p=0.063 | 1.474/p=0.208 | 2.113/p=0.077 | 3.404/p=0.009 | 4.171/p=0.002 | 1.515/p=0.195 | 3.809 p=0.004 |
| | 2 vs 5<0.05 | | | | 2 vs 3<0.05 | 1 vs 5<0.05 | | 2 vs 5<0.05 |
| | | | | | 2 vs 4<0.05 | 2 vs 4<0.05 | | |
| | | | | | 2 vs 5<0.05 | 2 vs 5<0.05 | | |

TABLE 3: Relationship between HLBCS and sociodemographic characteristics (continued).

| | Nutrition | Physical activity | Hygiene | Sleep | Environmental cleanliness | Health responsibility | Addiction | Total |
|--|-------------------------------------|-------------------------------------|-------------------------------------|----------------|--|-------------------------------------|-----------------------------------|------------------------------------|
| Father's educational status | | | | | | | | |
| Illiterate | 47.52±6.03 | 9.62±1.92 | 22.41±2.10 | 4.90±1.13 | 16.75±1.80 | 24.37±2.99 | 13.09±2.00 | 138.70±12.96 |
| Primary school | 47.44±5.85 | 9.83±1.85 | 22.62±1.94 | 4.85±1.02 | 16.48±2.11 | 24.46±2.93 | 13.20±1.89 | 138.91±12.70 |
| Middle school | 47.50±5.57 | 9.94±1.86 | 22.71±1.76 | 4.91±1.05 | 16.72±1.85 | 24.79±2.56 | 13.38±1.80 | 139.98±12.24 |
| High school | 48.79±5.49 | 10.25±1.76 | 22.93±1.62 | 4.97±1.04 | 16.91±1.43 | 24.92±2.35 | 13.41±1.73 | 142.20±11.09 |
| University | 48.69±5.49 | 10.15±1.85 | 22.63±1.80 | 4.95±1.05 | 16.90±1.63 | 24.79±2.75 | 13.49±1.85 | 141.63±12.62 |
| Anova/p value | 3.640/p=0.006 | 3.101/p=0.015 | 2.135/p=0.074 | 0.395/p=0.812 | 2.224/p=0.064 | 1.193/p=0.312 | 1.272/p=0.279 | 3.321 p=0.010 |
| Mother's profession | | | | | | | | |
| Housewife | 48.20±5.54 | 10.07±1.82 | 22.76±1.72 | 4.94±1.04 | 16.80±1.73 | 24.75±2.58 | 13.45±1.73 | 141.00±11.77 |
| Officer | 48.67±6.04 | 10.08±1.91 | 22.64±1.93 | 4.89±1.11 | 16.91±1.64 | 24.68±2.85 | 13.32±2.00 | 141.22±13.83 |
| Employee | 48.16±5.53 | 10.02±1.84 | 22.70±1.79 | 4.95±1.02 | 16.80±1.67 | 24.90±2.64 | 13.41±1.81 | 140.98±11.98 |
| Other occupations and death | 47.39±4.25 | 10.14±1.71 | 22.53±1.73 | 4.95±0.94 | 16.31±2.04 | 24.26±2.53 | 12.70±2.01 | 138.31±9.99 |
| Anova/p value | 0.859/p=0.462 | 0.105/p=0.957 | 0.454/p=0.714 | 0.219/p=0.883 | 1.446/p=0.228 | 0.876/p=0.453 | 2.269/p=0.079 | 0.683 p=0.562 |
| Father's profession | | | | | | | | |
| Not working | 47.72±5.70 | 9.63±2.06 | 22.44±2.77 | 4.94±1.05 | 16.65±2.20 | 24.07±3.36 | 13.17±1.83 | 138.65±14.65 |
| Officer | 48.65±5.88 | 10.06±1.87 | 22.64±1.84 | 4.90±1.10 | 16.85±1.74 | 24.71±2.79 | 13.41±1.96 | 141.25±13.50 |
| Employee | 48.16±5.55 | 10.07±1.81 | 22.77±1.66 | 4.95±1.03 | 16.86±1.60 | 24.85±2.53 | 13.43±1.76 | 141.12±11.51 |
| Other occupations and death | 47.95±4.73 | 10.44±1.74 | 22.77±1.53 | 4.93±0.95 | 16.22±1.98 | 24.64±2.40 | 13.01±1.76 | 140.00±10.34 |
| Anova/p value | 0.927/p=0.427 | 2.292/p=0.077 | 0.942/p=0.420 | 0.205/p=0.893 | 3.390 p=0.017 2 vs 4<0.05 3 vs 4<0.05 | 1.966/p=0.117 | 1.569/p=0.195 | 1.087 p=0.353 |
| Economic level of the family | | | | | | | | |
| Income is less than expenses | 48.87±5.42 | 10.16±1.85 | 22.85±1.72 | 4.95±1.06 | 16.90±1.65 | 24.95±2.62 | 13.51±1.76 | 142.23±12.00 |
| Income is equal to expenses | 47.26±5.78 | 9.88±1.81 | 22.50±1.83 | 4.90±1.03 | 16.63±1.75 | 24.46±2.61 | 13.16±1.91 | 138.83±12.25 |
| Income is more than expenses | 46.75±5.57 | 10.20±2.02 | 22.24±2.41 | 4.89±0.97 | 16.96±2.32 | 23.82±3.56 | 13.27±1.77 | 138.17±13.53 |
| Anova/p value | 12.78/p=0.000 1 vs 2<0.05 | 3.320/p=0.036 1 vs 2<0.05 | 6.178/p=0.002 1 vs 2<0.05 | 0.348/p=0.706 | 3.754/p=0.024 1 vs 2<0.05 | 6.590/p=0.001 1 vs 2<0.05 | 5.148 p=0.006 1vs2<0.05 | 11.698 p=0.000 1vs2<0.05 |
| Overweight level according to BMI | | | | | | | | |
| <5 (Low weight) | 48.35±5.48 | 9.95±1.90 | 22.72±1.87 | 5.00±0.93 | 16.87±1.41 | 24.91±2.31 | 13.62±1.52 | 141.45±11.31 |
| 5-85 (Normal weight) | 48.35±5.71 | 10.12±1.84 | 22.74±1.73 | 4.95±1.04 | 16.82±1.68 | 24.83±2.54 | 13.39±1.84 | 141.23±12.22 |
| 85-95 (Overweight) | 48.61±5.32 | 10.11±1.70 | 22.74±1.77 | 4.95±1.16 | 16.72±2.01 | 24.60±2.90 | 13.43±1.90 | 141.19±12.32 |
| 95> (Obese) | 47.03±5.12 | 9.66±1.92 | 22.46±2.14 | 4.74±1.05 | 16.71±1.77 | 24.24±3.44 | 13.03±1.83 | 137.91±12.85 |
| Anova/p value | 1.961/p=0.118 | 2.160/p=0.091 | 0.775/p=0.508 | 1.362/p=0.253 | 0.297/p=0.827 | 1.835/p=0.139 | 1.987/p=0.114 | 2.438 p=0.063 |
| Is there an overweight member in the family? | | | | | | | | |
| Yes | 47.56±5.53 | 9.98±1.87 | 22.57±1.86 | 4.78±1.07 | 16.66±1.87 | 24.46±2.80 | 13.28±1.82 | 139.34±12.21 |
| No | 48.82±5.61 | 10.1±1.82 | 22.83±1.72 | 5.05±1.01 | 16.92±1.56 | 24.99±2.51 | 13.47±1.82 | 142.25±12.10 |
| t-test/p value | -3.936/p=0.000 | -1.447/p=0.148 | -2.450/p=0.014 | -4.540/p=0.000 | -2.615/p=0.009 | -3.469/p=0.001 | -1.771 p=0.077 | -4.187 p=0.000 |

Anova: Analysis of variance; BMI: Body mass index

TABLE 4: Predictors affecting healthy lifestyle behaviors in primary school students

| Independent variables | B | SE | Beta (β) | t value | p value | F | Model (p) | R ² | Durbin Watson |
|---|---------|-------|------------------|---------|---------|-------|-----------|----------------|---------------|
| Constant | 135.235 | 1.233 | - | 109.710 | 0.000* | 9.217 | 0.000* | 0.078 | 1.813 |
| Gender (female) | 3.791 | 0.858 | 0.155 | 4.416 | 0.000* | | | | |
| Age (8 age) | -0.368 | 1.021 | -0.014 | -0.361 | 0.719 | | | | |
| Grade (3 rd) | 3.405 | 1.047 | 0.123 | 3.253 | 0.001* | | | | |
| Is there an overweight member in the family? (no) | -1.684 | 0.862 | -0.068 | -1.954 | 0.051 | | | | |
| Mother education (university and above) | 3.669 | 1.152 | 0.143 | 3.187 | 0.001* | | | | |
| Father education (university and above) | 1.556 | 1.141 | 0.061 | 1.364 | 0.173 | | | | |
| Economic status (Income is more than expenses) | 3.416 | 0.887 | 0.135 | 3.849 | 0.000* | | | | |

*p<0.05. SE: standard error of coefficient; β : standardized regression coefficient; R²: proportion of variation in dependent variable explained by regression model; p: the level of statistical significance.

of the regression analysis, gender was determined as an important variable affecting healthy lifestyle behaviors ($\beta=0.155$). It was determined that girls had higher lifestyle behaviors than boys. Xiang et al. also found that girls were more sensitive to healthy lifestyle factors than boys.²⁴ Temircan found that the healthy lifestyle behavior of female school-aged children's was higher than that of male school-aged children's.²⁵ Our research finding and the findings of previous studies show that girls have more positive lifestyle behaviors than boys.

In this study, mother's education level ($\beta=0.143$) was the other important variable affecting healthy lifestyle behaviors. While the father's education level was found to be associated with healthy lifestyle behaviors in univariate analysis, it was not found to be a predictor in multiple linear regression analysis. Children with higher educational level of their parents had higher healthy lifestyle behaviors. In the study of Musić Milanović et al. low parental education level was found to be a risk factor for not exercising.¹⁵ Previous studies on the subject in adolescents show that parental education level is an important predictor of children's healthy lifestyle behaviors.^{26,27} Our research finding is similar to previous research findings and supports the literature. Higher educated parents are typically more aware of and knowledgeable about health-related issues. These parents can provide more information and guidance to their children on topics such as healthy eating, regular exercise, hygiene and avoiding harmful habits. Parents are often important role models for their chil-

dren. When parents with higher levels of education demonstrate their own healthy lifestyle behaviors, they can set a positive example for their children. Educated parents can also allocate more resources to healthy lifestyle behaviors. For example, factors such as access to healthy foods, sports and physical activity opportunities, and access to health services can support children's healthy lifestyle behaviors.^{28,29}

The other important variable affecting healthy lifestyle behaviors in our study was the income status of children ($\beta=0.135$). The perception of the child's income level revealed that children with good income status had higher healthy lifestyle behaviors. In the study of Xiang et al. it was determined that children from low-medium family wealth had lower levels of physical activity than children from high family wealth.²⁴ Although not in primary school-aged children's, it shows that low-income perception is an important predictor affecting healthy lifestyle behaviors in adolescents.^{27,30} Our research finding and previous research findings suggest that income status is an important predictor of healthy lifestyle behaviors. Families with better income generally have access to healthier foods and can provide their children with a balanced diet from a variety of food groups. They may also invest more in physical activity opportunities, such as gym memberships and opportunities to join sports teams. This can encourage children to engage in regular physical activity.^{12,15}

In our study, another important factor affecting the healthy lifestyle behaviors of primary school-aged

children's was the grade level of the school-aged children's. Younger school-aged children's in the 3rd grade ($\beta=0.123$) had more positive lifestyle behaviors than 4th graders. Depboylu and Kaner reported that the nutritional compliance of primary school children with healthy lifestyle behaviors was significantly higher than that of middle school children.³¹ In the literature, younger age is associated with more dietary compliance, regular exercise, less screen time, and enough sleep length.^{21,31,32} There are studies showing that unhealthy dietary practices, inadequate physical activity, tobacco use, alcohol consumption and disturbances in sleep patterns increase with age.^{21,33} Considering that healthy lifestyle behaviors become negative and risky behaviors increase with increasing age, it is of great importance to gain healthy lifestyle behaviors in early childhood.

The school health nurse plays a crucial role in promoting healthy lifestyle behaviors in primary school-aged children's. This role is critical for children to acquire healthy habits and adopt long-term healthy lifestyles. School health nurses can create health education programs and provide trainings on topics such as healthy nutrition, sleep, exercise and hygiene. They can also help school-aged children's and their families achieve their personal health goals by providing individual counselling on healthy lifestyles. School health nurses can provide early diagnosis and treatment by conducting health screenings such as eye, dental and hearing screenings. They can encourage children to lead an active lifestyle by organizing sports activities and games. They can help school-aged children's cope with stress, anxiety and other emotional problems and ensure that they receive professional help when necessary. They ensure that the school environment is clean and hygienic and educate school-aged children's about personal hygiene. In this way, the school health nurse protects both the physical and psychological health of school-aged children's, supports their general well-being and helps them adopt healthy lifestyle behaviors.^{17,34,35}

STRENGTHS AND LIMITATIONS

The healthy lifestyle behaviors of primary school children and the sociodemographic factors influenc-

ing these practices were both thoroughly examined in our study. A further advantage of the research is that it was performed with a large sample ($n=1,240$). Our study also has limitations. In the study, healthy lifestyle behaviors were associated with sociodemographic variables, but the relationship with environmental and cultural variables within the scope of social determinants of health was not evaluated. In future studies, addressing the social determinants of health in detail may yield stronger results.

CONCLUSION

It was shown in this study that school-aged children's in primary schools exhibited a good level of healthy lifestyle behaviors. The mother's degree of education, income level, class level, and gender were the most significant determinants of healthy lifestyle behaviors. In line with these results, it is recommended to develop policies to improve sociodemographic determinants that affect healthy lifestyle behaviors. These may include increasing the education level of parents and providing social assistance to low-income families. According to these findings, it is recommended to create and support programs such as nutrition and physical activity in schools for healthy lifestyle behaviors among primary school-aged children's who are disadvantaged in terms of sociodemographic variables. In addition, the content of the programs to be planned should be organized according to demographic differences such as grade level and gender. School health nurses should plan and implement interventions for healthy lifestyle behaviors in line with the results of this study.

IMPLICATIONS FOR PRACTICE

Although the development of healthy lifestyle behaviors during childhood is important for the permanence of these behaviors, there is little evidence on children's healthy lifestyle behaviors. Starting from primary school, school health nurses should assess children's healthy lifestyle behaviors and the predictors affecting them. In line with these assessments, they should help children develop healthy lifestyle behaviors by taking individual differences into consideration.

Acknowledgments

The authors thanks to primary school children who participate in study.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family mem-

bers of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Asiye Kartal, Eda Kılınç İşleyen, Edanur Özkaya Bozkurt; **Design:** Asiye Kartal, Eda Kılınç İşleyen, Edanur Özkaya Bozkurt; **Control/Supervision:** Asiye Kartal; **Data Collection and/or Processing:** Eda Kılınç İşleyen, Edanur Özkaya Bozkurt; **Analysis and/or Interpretation:** Eda Kılınç İşleyen, Edanur Özkaya Bozkurt; **Literature Review:** Asiye Kartal, Eda Kılınç İşleyen, Edanur Özkaya Bozkurt; **Writing the Article:** Edanur Özkaya Bozkurt, Eda Kılınç İşleyen; **Critical Review:** Asiye Kartal.

REFERENCES

- Marques A, Bordado J, Tesler R, Demetriou Y, Sturm DJ, de Matos MG. A composite measure of healthy lifestyle: a study from 38 countries and regions from Europe and North America, from the Health Behavior in School-Aged Children survey. *Am J Hum Biol.* 2020;32(6):e23419. [Crossref] [PubMed]
- WHO [Internet]. A healthy lifestyle-WHO recommendations. © 2010 WHO [Cited: January 13, 2024] Available from: [Link]
- Marques A, Loureiro N, Avelar-Rosa B, Naia A, Matos MG. Adolescents' healthy lifestyle. *J Pediatr (Rio J).* 2020;96(2):217-24. [Crossref] [PubMed] [PMC]
- Owen KB, Parker PD, Astell-Burt T, Lonsdale C. Regular physical activity and educational outcomes in youth: a longitudinal study. *J Adolesc Health.* 2018;62(3):334-40. [Crossref] [PubMed]
- Ng R, Sutradhar R, Yao Z, Wodchis WP, Rosella LC. Smoking, drinking, diet and physical activity-modifiable lifestyle risk factors and their associations with age to first chronic disease. *Int J Epidemiol.* 2020;49(1):113-30. [Crossref] [PubMed] [PMC]
- Scaglioni S, De Cosmi V, Ciappolino V, Parazzini F, Brambilla P, Agostoni C. Factors Influencing Children's Eating Behaviours. *Nutrients.* 2018;10(6):706. [Crossref] [PubMed] [PMC]
- Stevenson D. Settings for Health Promotion. In: Snelling A, ed. *Introduction to Health Promotion.* 2nd ed. New Jersey: John Wiley&Sons. 2024. p.253-77.
- Pagerols M, Prat R, Rivas C, Español-Martin G, Puigbó J, Pagespetit È, et al. The impact of psychopathology on academic performance in school-age children and adolescents. *Sci Rep.* 2022;12(1):4291. [Crossref] [PubMed] [PMC]
- UNESCO. Ready to learn and thrive: school health and nutrition around the world. [Cited: December 12, 2023]. [Link]
- Puia A, Leucuta DC. Children's lifestyle behaviors in relation to anthropometric indices: a family practice study. *Clujul Med.* 2017;90(4):385-91. [Crossref] [PubMed] [PMC]
- D'Souza NJ, Kuswara K, Zheng M, Leech R, Downing KL, Lioret S, et al. A systematic review of lifestyle patterns and their association with adiposity in children aged 5-12 years. *Obes Rev.* 2020;21(8):e13029. [Crossref] [PubMed]
- Whiting S, Buoncristiano M, Gelius P, Abu-Omar K, Pattison M, Hyska J, et al. Physical activity, screen time, and sleep duration of children aged 6-9 years in 25 countries: an analysis within the WHO European childhood obesity surveillance initiative (COSI) 2015-2017. *Obes Facts.* 2021;14(1):32-44. [Crossref] [PubMed] [PMC]
- Lekše R, Godec D, Prosen M. Determining the impact of lifestyle on the health of primary school children in slovenia through mixed membership focus groups. *J Community Health.* 2023;48(5):857-69. [Crossref] [PubMed] [PMC]
- Mulitauopele C, Israel W, Rosenberg S, Brooks-Russell A, Ma M, Dinger MK. The association of school learning environment with adolescent health behaviors. *J Sch Health.* 2023;93(8):649-58. [Crossref] [PubMed]
- Musić Milanović S, Buoncristiano M, Krizan H, Rathmes G, Williams J, Hyska J, et al. Socioeconomic disparities in physical activity, sedentary behavior and sleep patterns among 6- to 9-year-old children from 24 countries in the WHO European region. *Obes Rev.* 2021;22 Suppl 6:e13209. [Crossref] [PubMed]
- Rawal T, van Schayck OCP, Willeboordse M, Arora M, Bhaumik S, Bhagra A, et al. How to promote a healthy lifestyle among schoolchildren: development of an intervention module (i-PROMISE). *Public Health Pract (Oxf).* 2022;3:100262. [Crossref] [PubMed] [PMC]
- Kostenius C. School nurses' experiences with health dialogues: a Swedish case. *J Sch Nurs.* 2023;39(5):345-56. [Crossref] [PubMed] [PMC]
- Öcal EE, Önsüz MF. Reliability and validity of healthy lifestyle behaviours scale in children. *European Journal of Public Health.* 2020;30(Supplement_5). [Crossref]
- George D, Mallery M. *SPSS for Windows Step by Step: A Simple Guide and Reference.* 17.0 update. 10th ed. Boston: Pearson; 2010.
- Neyzi O, Günöz H, Furman A, Bundak R, Gökçay G, Darendeliler F, et al. Türk çocuklarında vücut ağırlığı, boy uzunluğu, baş çevresi ve vücut kitle indeksi referans değerleri. *Çocuk Sağlığı ve Hastalıkları Dergisi.* 2008;51(1):1-14. [Link]
- Hormenu T. Dietary intake and its associated factors among in-school adolescents in Ghana. *PLoS One.* 2022;17(5):e0268319. [Crossref] [PubMed] [PMC]
- López-Bueno R, López-Sánchez GF, Casajús JA, Calatayud J, Tully MA, Smith L. Potential health-related behaviors for pre-school and school-aged children during COVID-19 lockdown: a narrative review. *Prev Med.* 2021;143:106349. [Crossref] [PubMed] [PMC]
- Day RE, Sahota P, Christian MS. Effective implementation of primary school-based healthy lifestyle programmes: a qualitative study of views of school staff. *BMC Public Health.* 2019;19(1):1239. [Crossref] [PubMed] [PMC]
- Xiang H, Feng X, Lin L, Luo S, Liu X, Chen D, et al. Association between healthy lifestyle factors and health-related quality of life among Chinese adolescents: the moderating role of gender. *Health Qual Life Outcomes.* 2023;21(1):119. [Crossref] [PubMed] [PMC]

25. Temircan Z. Ergenlerin sağlıklı yaşam biçimi davranışları ile okul aidiyeti arasındaki ilişkinin incelenmesi [Investigation of the relationship between adolescents' healthy lifestyle behaviors and school belonging]. *Journal of Mehmet Akif Ersoy University Institute of Educational Sciences*. 2022;10(13):71-87. [\[Link\]](#)
26. Ardic A, Esin MN. Factors associated with healthy lifestyle behaviors in a sample of Turkish adolescents: a school-based study. *J Transcult Nurs*. 2016;27(6):583-92. [\[Crossref\]](#) [\[PubMed\]](#)
27. Daşkan Z. Healthy lifestyle behaviors and related factors in high school adolescents: a cross-sectional study (Izmir/Turkey). *Vulnerable Children and Youth Studies*. 2023;18(2):156-68. [\[Crossref\]](#)
28. Chen L, Hong J, Xiong D, Zhang L, Li Y, Huang S, et al. Are parents' education levels associated with either their oral health knowledge or their children's oral health behaviors? A survey of 8446 families in Wuhan. *BMC Oral Health*. 2020;20(1):203. [\[Crossref\]](#) [\[PubMed\]](#) [\[PMC\]](#)
29. de Buhr E, Tannen A. Parental health literacy and health knowledge, behaviours and outcomes in children: a cross-sectional survey. *BMC Public Health*. 2020;20(1):1096. [\[Crossref\]](#) [\[PubMed\]](#) [\[PMC\]](#)
30. Gur K, Kılınç E, Yayıkci E, Değer K, Tekin E. Examination of predictive factors healthy lifestyle behaviours and compare coping styles with stress of adolescents with and without hearing loss: a comparative study. *Clinical and Experimental Health Sciences*. 2022;12(4):875-84. [\[Link\]](#)
31. Depboyu GY, Kaner G. Younger age, higher father education level, and healthy lifestyle behaviors are associated with higher adherence to the Mediterranean diet in school-aged children. *Nutrition*. 2023;114:112166. [\[Crossref\]](#) [\[PubMed\]](#)
32. Lioret S, Campbell KJ, McNaughton SA, Cameron AJ, Salmon J, Abbott G, et al. Lifestyle patterns begin in early childhood, persist and are socioeconomically patterned, confirming the importance of early life interventions. *Nutrients*. 2020;12(3):724. [\[Crossref\]](#) [\[PubMed\]](#) [\[PMC\]](#)
33. Jonsson KR, Corell M, Löfstedt P, Adjei NK. The clustering of multiple health and lifestyle behaviors among Swedish adolescents: a person-oriented analysis. *Front Public Health*. 2023;11:1178353. [\[Crossref\]](#) [\[PubMed\]](#) [\[PMC\]](#)
34. Silva AA, Gubert FDA, Barbosa Filho VC, Freitas RWJF, Vieira-Meyer APGF, Pinheiro MTM, et al. Health promotion actions in the School Health Program in Ceará: nursing contributions. *Rev Bras Enferm*. 2021;74(1):e20190769. English, Portuguese. [\[Crossref\]](#) [\[PubMed\]](#)
35. Dolgun E, Yavuz van Giersbergen M, Durmaz Akyol A. Healthy life style behaviors changes of the nursing students: 2004-2014. *International Journal of Caring Sciences*. 2020;13(2):1155-61. [\[Link\]](#)